

Key IYCF-E Indicators

IYCF-E Indicators - to be included in monthly data collection and reporting
of breastfeeding support groups formed (including MTMSG and Care Groups)
of pregnant and lactating women attending breastfeeding support groups (disaggregated by pregnant and lactating)
of pregnant and lactating women referred for individual counselling (disaggregated by pregnant/lactating)
of pregnant and lactating women who received individual counselling (disaggregated by pregnant/lactating)
of infants 0-6 months who need BMS, after assessment by a qualified health workers
of infants 0-6 months who received BMS, after assessment by a qualified health worker
of reports of BMS code violations (disaggregated by type of violation)
of safe breastfeeding areas established
of pregnant and lactating women attending safe breastfeeding areas
of health workers trained on breastfeeding counselling (assessment of mother baby pair and counselling skills)
of health workers trained on BMS Code and Reporting Form for Code Violations

IYCF Indicators - to be included in bi-annual surveys (SMART, MICS, IYCF KAP etc.)
of children 0-23 months who are fed with feeding bottles/teats
of CMAM structures with a satisfactory IYCF-E integration score (quarterly joint supervision visit)
Early initiation; proportion of children 0-23 months who were put to the breast within 30 minutes of birth
Exclusive breastfeeding; proportion of infants 0-5 months of age who are fed exclusively with breast milk in the previous 24 hrs
Continued breastfeeding at 1 year of age; proportion of children 12-15 months of age who are fed breast milk in the previous 24 hrs
Continued breastfeeding up to 2 years of age; proportion of children 20-23 months of age who are fed breast milk in the previous 24 hrs
Timely introduction of complementary feeding; proportion of infants 6-9 months who receive breastmilk and a solid or semi-solid in the previous 24 hrs
Minimum dietary diversity; proportion of children 6-23 months who received food from 4 or more food groups in the previous 24 hrs
Minimum meal frequency; proportion of breastfed and non-breastfed children 6-23 months who received solid, semi-solid, or soft foods the minimum number of times or more